

Covariant Description of the Inhomogeneous Mixmaster Chaos

R. Benini^{12†} and G. Montani^{23‡}

¹*Dipartimento di Fisica - Università di Bologna and INFN
Sezione di Bologna, via Irnerio 46, 40126 Bologna, Italy*

²*ICRA—International Center for Relativistic Astrophysics c/o Dipartimento di Fisica (G9)
Università di Roma “La Sapienza”, Piazza A.Moro 5 00185 Roma, Italy*

³*ENEA C.R. Frascati (U.T.S. Fusione),
via Enrico Fermi 45, 00044 Frascati, Rome, Italy*

[†]*riccardo.benini@icra.it*

[‡]*montani@icra.it*

We outline the covariant nature of the chaos characterizing the generic cosmological solution near the initial singularity. Our analysis is based on a “gauge” independent ADM-reduction of the dynamics to the physical degrees of freedom, and shows that the dynamics is isomorphic point by point in space to a billiard on a Lobachevsky plane. The Jacobi metric associated to the geodesic flow is constructed and a non-zero Lyapunov exponent is explicitly calculated. The chaos covariance emerges from the independence of the form of the lapse function and the shift vector.

In recent years, the chaoticity of the homogeneous Mixmaster model has been widely studied in the literature (see^{1–4}) in view of understanding the features of its covariant nature. Two convincing arguments, appeared in,^{1,4} support the idea that the Mixmaster chaos (described by the invariant measure introduced in^{3,5}) remains valid in any system of coordinates.

The main issue of the present work is to show that the property of space-time covariance can be extended to the inhomogeneous Mixmaster model.

A generic cosmological solution is represented by a gravitational field having available all its degrees of freedom and, therefore, allowing to specify a generic Cauchy problem. In the Arnowitt-Deser-Misner (ADM) formalism, the metric tensor corresponding to such a generic model takes the form

$$d\Gamma^2 = N^2 dt^2 - \gamma_{\alpha\beta} (dx^\alpha + N^\alpha dt)(dx^\beta + N^\beta dt) \quad (1)$$

where N and N^α denote (respectively) the lapse function and the shift-vector, $\gamma_{\alpha\beta}$ the 3-metric tensor of the spatial hyper-surfaces Σ^3 for which $t = \text{const}$, being⁶

$$\gamma_{\alpha\beta} = e^{q_a} \delta_{ad} O_b^a O_c^d \partial_\alpha y^b \partial_\beta y^c \quad a, b, c, d, \alpha, \beta = 1, 2, 3; \quad (2)$$

$q^a = q^a(x, t)$ and $y^b = y^b(x, t)$ are six scalar functions, and $O_b^a = O_b^a(x)$ a $SO(3)$ matrix. By the metric tensor (2), the action for the gravitational field is

$$S = \int_{\Sigma^{(3)} \times \mathbb{R}} dt d^3x (p_a \partial_t q^a + \Pi_d \partial_t y^d - NH - N^\alpha H_\alpha), \quad (3)$$

$$H = \frac{1}{\sqrt{\gamma}} \left[\sum_a (p_a)^2 - \frac{1}{2} \left(\sum_b p_b \right)^2 - \gamma^{(3)} R \right] \quad (4)$$

$$H_\alpha = \Pi_c \partial_\alpha y^c + p_a \partial_\alpha q^a + 2p_a (O^{-1})_a^b \partial_\alpha O_b^a; \quad (5)$$

in (4) and (5) p_a and Π_d are the conjugate momenta to the variables q^a and y^b respectively, and the $^{(3)}R$ is the Ricci 3-scalar which plays the role of a potential term. We use the Hamiltonian constraints $H = H_\alpha = 0$ to reduce the dynamics to the physical degrees of freedom; the super-momentum constraints can be diagonalized and explicitly solved by choosing the function y^a as special coordinates:

$$\Pi_b = -p_a \frac{\partial q^a}{\partial y^b} - 2p_a (O^{-1})_a^c \frac{\partial O_c^a}{\partial y^b}. \quad (6)$$

$$S = \int_{\Sigma^{(3)} \times \mathbb{R}} d\eta d^3y \left(p_a \partial_\eta q^a + 2p_a (O^{-1})_a^c \partial_\eta O_c^a - NH \right). \quad (7)$$

The potential term appearing in the super-Hamiltonian behaves as an infinite potential wall as the determinant of the 3-metric goes to zero, and we can model it as follows

$$U = \sqrt{\gamma}^{(3)} R = \sum_a \Theta(Q_a) \quad \Theta(x) = \begin{cases} +\infty & \text{if } x < 0, \\ 0 & \text{if } x > 0. \end{cases} \quad (8)$$

where the Q_a 's are called *anisotropy parameters*. By (8) the Universe dynamics evolves independently in each space point and the point-Universe can move within a dynamically-closed domain Γ_Q only⁶ (see figure (1)). Since in Γ_Q the potential U asymptotically vanishes, near the singularity we have $\partial p_a / \partial \eta = 0$; then the term $2p_a (O^{-1})_a^c \partial_\eta O_c^a$ in (7) behaves as an exact time-derivative that can be ruled out of the variational principle.

The ADM reduction is completed by introducing the so-called Misner-Chitré like (τ, ξ, θ) variables,¹ in terms of which the anisotropy parameters Q_a become τ independent: When expressed in terms of such variables the super-Hamiltonian con-

$$\begin{aligned} Q_1 &= \frac{1}{3} - \frac{\sqrt{\xi^2 - 1}}{3\xi} (\cos \theta + \sqrt{3} \sin \theta) \\ Q_2 &= \frac{1}{3} - \frac{\sqrt{\xi^2 - 1}}{3\xi} (\cos \theta - \sqrt{3} \sin \theta) \\ Q_3 &= \frac{1}{3} + \frac{2\sqrt{\xi^2 - 1}}{3\xi} \cos \theta \end{aligned} \quad (9)$$




Fig. 1. $\Gamma_Q(\xi, \theta)$

straint can be solved in the domain Γ_Q according to the ADM prescription:

$$-p_\tau \equiv \epsilon = \sqrt{(\xi^2 - 1)p_\xi^2 + \frac{p_\theta^2}{\xi^2 - 1}} \quad (10)$$

$$\delta S_{\Gamma_Q} = \delta \int d\eta d^3y (p_\xi \partial_\eta \xi + p_\theta \partial_\eta \theta - \epsilon \partial_\eta \tau) = 0. \quad (11)$$

By virtue of the asymptotic limit (8) and the Hamilton equations associated with (11) it follows that ϵ is a constant of motion, i.e. $d\epsilon/d\eta = \partial\epsilon/\partial\eta = 0 \Rightarrow \epsilon = E(y^a)$; furthermore $\epsilon\partial_\eta\tau$ behaves as an exact time derivative.

This dynamical scheme allows us to construct the Jacobi metric corresponding to the dynamical flow, and the line element reads

$$ds^2 = E^2(y^a) \left(\frac{d\xi^2}{(\xi^2 - 1)} + (\xi^2 - 1)d\theta^2 \right). \quad (12)$$

Here the space coordinates behaves like external parameters since the evolution is spatially uncorrelated. The Ricci scalar takes the value $R = -2/E^2$, so that (12) describes a two-dimensional Lobachevsky space; the role of the potential wall (8) consists of cutting a closed domain Γ_Q on such a negative curved surface.

In this scheme the Lyapunov exponent associated to the geodesic deviation along the direction orthonormal to the geodesic 4-velocity can be evaluated, and it results to be equal to

$$\lambda(y^a) = \frac{1}{\epsilon(y^a)} > 0. \quad (13)$$

Hence the chaotic behavior of the inhomogeneous Mixmaster model can be described in a generic gauge (*i.e.* without assigning the form of the lapse function and of the shift vector) as soon as Misner-Chitré like variables are adopted. The value (13) is a positive definite function of the space point, making this calculation extendible point by point to the whole space.

References

1. G. Imponente and G. Montani, Phys. Rev. D **63** (2001) 103501
2. G. Montani, Class. Quant. Grav. **18** (2001) 193
3. A. A. Kirillov and G. Montani, Phys. Rev. D **56** (1997) 6225.
4. N. J. Cornish and J. J. Levin, Phys. Rev. Lett. **78**, 998 (1997)
5. D.F. Chernoff and J.D. Barrow, Phys. Rev. Lett, **50**, 134,(1983).
6. R. Benini, G. Montani and R. Benini, Phys. Rev. D **70** (2004) 103527